

Spacetime fluctuations in AdS/CFT

Erik Verlinde^a and Kathryn M. Zurek^b

^a*Institute of Physics, University of Amsterdam,
Amsterdam, The Netherlands*

^b*Walter Burke Institute for Theoretical Physics, California Institute of Technology,
Pasadena, CA, U.S.A.*

E-mail: e.p.verlinde@uva.nl, kzurek@caltech.edu

ABSTRACT: We compute fluctuations in the modular energy of the vacuum associated with a Rindler-wedge in AdS spacetime in the context of AdS/CFT. We discuss the possible effect of these energy fluctuations on the spacetime geometry, and on the traversal time of a light beam propagating from the boundary to the bulk and back.

KEYWORDS: AdS-CFT Correspondence, Models of Quantum Gravity

ARXIV EPRINT: [1911.02018](https://arxiv.org/abs/1911.02018)

Contents

1	Introduction	1
1.1	Energy fluctuations in the AdS/CFT vacuum state	2
1.2	Summary of the main results	3
2	The geometric setup	4
2.1	From Poincaré to AdS-Rindler coordinates	5
2.2	The regularized horizon area of AdS-Rindler and topological black holes	6
3	Fluctuations in the modular Hamiltonian	7
3.1	The modular Hamiltonian: bulk versus boundary	7
3.2	The gravitational replica method	8
3.3	Computation of the fluctuations in the modular Hamiltonian	11
4	Backreaction due to the fluctuating modular energy	12
4.1	The relation between the mass M and the modular hamiltonian K	13
4.2	Fluctuations in the Newtonian potential on the Rindler horizon	14
4.3	Horizon location and photon travel time	15
4.4	Fluctuations in horizon location and photon travel time	17
5	Discussion	19

1 Introduction

In a recent paper [1] we proposed that, generically, a theory of quantum gravity motivated by the holographic principle leads to fluctuations in the spacetime geometry that are measurable at macroscopic distances. Our reasoning was based on a number of steps. As a first step we argued that, when one considers a finite region of spacetime, the energy contained in that region, even in the vacuum, fluctuates. Specifically, we considered the fluctuations in a suitably defined energy, the so-called “modular Hamiltonian,” associated to a causal diamond, which describes the domain of causal dependence of a given (finite) spatial region.

The boundary of the causal diamond is locally described by a Rindler horizon, and thus carries a non-zero temperature and entropy, both of which are due to the quantum entanglement between the interior and exterior of the diamond. A crucial ingredient in the argumentation of [1] was that as a result the boundary of the causal diamond can be treated as a generalization of a black hole event horizon, in the sense that it obeys the same laws of (black hole) thermodynamics. From this assumption we showed that the size of the fluctuations in the modular Hamiltonian is directly determined by the amount of entanglement carried by the microscopic degrees of freedom associated with the diamond.

The second step of our reasoning was to argue that these energy fluctuations “gravitate,” meaning that they induce fluctuations in the spacetime geometry. In particular, it was shown that, due to the gravitational back-reaction, the modular energy fluctuations induce fluctuations in the location of the horizon at the edge of the causal diamond. The third step was to demonstrate that these fluctuations give rise to observable fluctuations in the phase of light between two mirrors of an interferometer that stretches from the center to the edge of the diamond.

The goal of this paper is to provide further evidence for the arguments and results presented in our work [1] by placing and repeating our calculation in the context of AdS/CFT. In the setting of AdS/CFT the microscopic description is well understood, which means that the various steps in our reasoning can be explicitly verified without uncertainty about the underlying microscopics. This allows us to make use of the many results obtained in the extensive literature on this subject to verify the central arguments and explicitly check the conclusions of our previous work.

1.1 Energy fluctuations in the AdS/CFT vacuum state

Empty AdS space corresponds to the ground state of the microscopic quantum theory, and is dual to the conformally invariant ground state $|\text{vac}\rangle$ of the CFT. Conformal transformations of the CFT are in one-to-one correspondence with isometries of the AdS geometry. The associated conserved charges H_ξ are given in terms of the stress energy tensor T_{ab}^{CFT} of the boundary CFT by

$$H_\xi = \int T_{ab}^{\text{CFT}} \xi^a dV^b, \quad (1.1)$$

where the infinitesimal volume element for the boundary CFT is denoted by dV^b , and ξ^a represents the conformal Killing vector on the boundary. While H_ξ is defined in terms of the boundary CFT, it also represents the conserved charge associated with the corresponding isometry of the bulk AdS-spacetime.

All the charges H_ξ annihilate the vacuum state $|\text{vac}\rangle$, which means that $H_\xi |\text{vac}\rangle = 0$. This fact implies that the (generalized) Hamiltonians H_ξ do not exhibit any vacuum fluctuations, since the matrix elements of H_ξ^2 are identically zero:

$$\langle \text{vac} | H_\xi^2 | \text{vac} \rangle = 0. \quad (1.2)$$

This does not mean, however, that there are no (local) energy fluctuations in the vacuum. Indeed, the stress tensor itself has a non-trivial two point function,

$$\langle \text{vac} | T_{ab}^{\text{CFT}}(x) T_{cd}^{\text{CFT}}(y) | \text{vac} \rangle \neq 0, \quad (1.3)$$

and thus fluctuates. The charges H_ξ are defined by integrating the currents $\xi^a T_{ab}^{\text{CFT}}$ over the entire boundary. If one would integrate over only part of the boundary, then the resulting quantities would no longer annihilate the vacuum, and will in general fluctuate.

In this paper we will consider the special Killing vector fields ξ_K that on the boundary generate conformal transformations that leave invariant a spherical co-dimension one surface and in the bulk correspond to boosts that leave invariant a Rindler horizon. These

vector fields naturally split the bulk as well as the boundary in to two parts, namely the two regions that are separated by the Rindler horizon in the bulk or a spherical co-dimension one surface on the boundary.

Let us denote the part of the boundary inside the spherical surface by B and its complement by \bar{B} . The conserved charge H_{ξ_K} can thus be decomposed as $H_{\xi_K} = K - \bar{K}$ where the quantity K is defined as

$$K = \int_B T_{ab}^{\text{CFT}} \xi_K^a dB^b, \quad (1.4)$$

and \bar{K} is given by an identical expression for \bar{B} . Here the integral is over the spherical volume B , such that dB^b is normal to this volume and hence points in the time direction. We will refer to the operator K as the modular energy or modular Hamiltonian. As an operator it does not annihilate the vacuum state: $K|\text{vac}\rangle \neq 0$ and hence it can fluctuate. In this paper we will explicitly compute these fluctuations using the dictionary of the AdS/CFT correspondence and indeed show that the result is non-zero,

$$\langle K^2 \rangle - \langle K \rangle^2 \neq 0. \quad (1.5)$$

It should again be emphasized that the non-zero value of these fluctuations are a consequence of the fact that we are considering a finite region of the AdS-spacetime.

1.2 Summary of the main results

The modular Hamiltonian K can, instead of by eq. (1.4), be alternatively defined in terms the density matrix σ describing the mixed quantum state of the CFT inside the region B . Schematically, σ is obtained by tracing over the Hilbert space $\mathcal{H}_{\bar{B}}$ for the outside region \bar{B} ,

$$\sigma = \text{tr}_{\mathcal{H}_{\bar{B}}} (|\text{vac}\rangle \langle \text{vac}|), \quad (1.6)$$

and in this case it can be shown that the density matrix takes a thermal form [2] in terms of the modular Hamiltonian,

$$\sigma = \frac{e^{-K}}{Z} \quad \text{with} \quad Z = \text{tr}(e^{-K}). \quad (1.7)$$

Note that the relation (1.7) defines K only up to a constant, since one can shift $K \rightarrow K + c$ and $Z \rightarrow e^{-c}Z$ without changing the state σ . We will use this freedom to identify the modular Hamiltonian with

$$K \equiv -\log \sigma. \quad (1.8)$$

With this choice the partition function Z in the vacuum state σ is simply equal to one.

The state σ represents a thermal state and hence it gives rise to thermal fluctuations. The first goal of this paper is to compute the thermal fluctuations in the expectation value of the operator K . This will be the starting point for determining the spacetime fluctuations. Normally, following the standard rules of quantum statistical physics, one computes the thermal fluctuations in the energy by computing the free energy

$$F_\beta = -\frac{1}{\beta} \log \text{tr} (e^{-\beta K}) \quad (1.9)$$

as a function of the inverse temperature β , and then takes the second derivative

$$\langle K^2 \rangle - \langle K \rangle^2 = -\frac{\partial^2}{\partial \beta^2} (\beta F_\beta) \Big|_{\beta=1}. \quad (1.10)$$

In the AdS/CFT context, however, it is not easy to compute F_β for arbitrary values of β . Instead, we will compute the free energy F_n only for integer values of $\beta = n$ using the replica method. Specifically, we will use the methods developed in the references [3–5].

First, it is clear from eq. (1.8) that the expectation value of the modular Hamiltonian K is equal to the entanglement entropy, which in a CFT that is dual to Einstein gravity can be computed using the Ryu-Takayanagi formula [6, 7]

$$\langle K \rangle = \frac{A(\Sigma)}{4G}. \quad (1.11)$$

Here $A(\Sigma)$ denotes the area of the extremal surface Σ in AdS that is anchored on the boundary on the edge of the region B . In the present context Σ also represents the Rindler horizon. A central result of this paper will be that the fluctuations in the modular Hamiltonian within AdS/CFT are given by

$$\langle K^2 \rangle - \langle K \rangle^2 = \frac{A(\Sigma)}{4G}. \quad (1.12)$$

This results holds in arbitrary dimensions for any strongly coupled CFT with a large N gravitational dual that is described by the Einstein action. We will subsequently study the implications of these modular energy fluctuations for the AdS-Rindler horizon and for the travel times of photons that probe the spacetime geometry near the horizon.

After completing our calculations, we became aware of earlier work with some overlap with the first part of our analysis. Our result (1.12) for the fluctuations in the modular Hamiltonian K agrees, after suitable identifications, with the expressions presented in ref. [8], which contains in addition to a bulk computation a CFT derivation of the temperature variation of the n -th Rényi entropy. The same physical quantity was considered in refs. [9], and [10], and was referred to as the capacity of entanglement. The emphasis of the present work is on the physical implications for the gravitational bulk physics rather than the boundary CFT. Since the concrete geometric form of the result eq. (1.12) differs from these earlier references, and to make the current analysis self-contained, we will present here our own analysis, which was performed independently.

2 The geometric setup

Consider a light signal that is first emitted from the boundary and then is reflected by a “mirror” at some location in the bulk, and is finally again received at a later moment on the boundary. The points of emission and reception on the boundary define a causal diamond that consists of all points that are in the causal future of the point of emission and the causal past of the point of reception. The past lightcone of the reception point intersects that future lightcone of the emission point along a co-dimension two surface, that represents a Ryu-Takayanagi (RT) surface Σ in the bulk. The causal diamond describes a wedge




Figure 1. Depiction of the causal diamond in AdS space anchored at the boundary. The red dotted line is the light signal from the boundary to a point in the bulk.

of Rindler-AdS space, with the Killing horizon located at its boundary. The light signal travels along this horizon and reflects at the point on the RT surface Σ . This situation is depicted in figure 1. Note that our geometric set-up is rather similar to some works seeking to derive the linearized Einstein equations from entanglement entropy [11, 12].

2.1 From Poincaré to AdS-Rindler coordinates

On the boundary we are thus dealing with a spherically symmetric causal diamond, whose causal domain extends in the bulk and describes one wedge of Rindler-AdS space. The Rindler horizon coincides with the RT surface Σ , which is anchored on the edge of the causal diamond on the boundary. We will work in two coordinate systems, one that describes the $(d+1)$ -dimensional AdS geometry near the boundary in terms of the familiar Poincaré coordinates:

$$ds^2 = L^2 \frac{dz^2 + dx_i^2 - dx_0^2}{z^2} \quad \text{with} \quad i = 1, \dots, d-1. \quad (2.1)$$

The finite spherical region on the boundary is described by $x_i^2 \leq R^2$. The shaded region in the bulk represents the spatial slice inside the causal diamond and is given by $z^2 + x_i^2 \leq R^2$. The full interior of the causal diamond is described by

$$R^2 - z^2 - x_i^2 + x_0^2 \geq 2R|x_0|. \quad (2.2)$$

It turns out that, while the region of interest is easily described in these coordinates, the calculation of the fluctuations in the modular Hamiltonian and the possible backreaction effects are more easily studied in a different coordinate system that is better adapted to the geometry of the causal diamond.

For this reason we introduce a second coordinate system that describes the interior of the causal diamond and puts it in the form of Rindler-AdS space (see for example ref. [2] for more details). The associated metric is

$$ds^2 = - \left(\frac{r^2}{L^2} - 1 \right) dt^2 + \left(\frac{r^2}{L^2} - 1 \right)^{-1} dr^2 + r^2 d\Sigma_{d-1}^2, \quad (2.3)$$

where

$$d\Sigma_{d-1}^2 = d\chi^2 + \sinh^2 \chi d\Omega_{d-2}^2 \quad (2.4)$$

describes the geometry of a hyperbolic plane. The radial coordinate obeys $r \geq L$, so that the coordinate t remains timelike. The boundary value $r = L$ corresponds to the location of the Rindler horizon. The relationship between the two coordinate systems is¹

$$\frac{R^2 - z^2 - x_i^2 + x_0^2}{2Rz} = \left(\frac{r^2}{L^2} - 1 \right)^{\frac{1}{2}} \cosh \frac{t}{L}, \quad \frac{x_0}{z} = \left(\frac{r^2}{L^2} - 1 \right)^{\frac{1}{2}} \sinh \frac{t}{L}. \quad (2.5)$$

The spatial x_i coordinates are replaced by the coordinate χ and the solid angles Ω by writing the norm $|x^i|$ as

$$|x_i| = R \tanh \chi. \quad (2.6)$$

One easily verifies that new coordinates fully cover the domain (2.2) and that the $r = L$ locus coincides with the boundary of this causal domain. The new (r, t, χ, Ω) -coordinate system thus describes only the interior of the causal diamond.

2.2 The regularized horizon area of AdS-Rindler and topological black holes

A central quantity in our calculations is the area of the AdS-Rindler horizon, which through the result of Ryu and Takayanagi measures the entanglement entropy between inside and outside of the causal diamond for the boundary CFT,

$$S = \frac{A(\Sigma)}{4G}. \quad (2.7)$$

The surface Σ is located at radius $r = L$. This means that we can write its area as

$$A(\Sigma) = L^{d-1} V_{d-1}, \quad (2.8)$$

where V_{d-1} denotes the dimensionless ‘volume’ of the $(d-1)$ -dimensional hyperbolic plane described by the metric eq. (2.4). The hyperbolic volume V_{d-1} is formally infinite, and hence needs to be regularized. We will therefore introduce a cut-off by restricting the range of the coordinate $\chi \leq \chi_c$. This leads to the following regularized expression

$$V_{d-1} = \Omega_{d-2} \int_0^{\chi_c} d\chi \sinh^{d-2} \chi, \quad (2.9)$$

where Ω_{d-2} is the area of a unit $(d-2)$ -sphere.

¹This coordinate transformation can be derived by making use of the embedding equations for AdS and tracing through the conformal mapping used in ref. [2] to go from Rindler space to the causal diamond.

A further important reason for introducing this regularization is that it allows us to consider geometries that generalize AdS-Rindler space by allowing the (modular) energy K to differ from its vacuum value. These geometries are called topological black holes and are described by the metric

$$ds^2 = -f(r)dt^2 + \frac{dr^2}{f(r)} + r^2 d\Sigma_{d-1}^2, \quad (2.10)$$

where the blackening factor $f(r)$ takes the form

$$f(r) = \frac{r^2}{L^2} - 1 - 2\Phi \frac{L^{d-2}}{r^{d-2}}, \quad (2.11)$$

obtained by solving the Einstein equations. The dimensionless quantity Φ is an integration constant, which physically represents the value of a Newtonian potential on the original location $r = L$ of the horizon. Φ can be written in terms of a mass M , which we will show below is the mass measured by the observer who uses t as their time coordinate. In this case, we will have

$$\Phi = \frac{8\pi GM}{(d-1)V_{d-1}L^{d-2}}, \quad (2.12)$$

where again V_{d-1} is the regularized hyperbolic volume of the horizon. A similar regularization is required to define the mass M , which without any regularization would be infinite. The ratio M/V_{d-1} is independent of the regularization and may be interpreted as the tension of a brane that lives at the location of the horizon. In fact, in the following we will make use of these geometries, with precisely this brane interpretation, to calculate the fluctuations in the values of the modular Hamiltonian.

3 Fluctuations in the modular Hamiltonian

Our goal is to compute the fluctuations in the modular Hamiltonian K in the vacuum state σ . These fluctuations are described by the two point function,

$$\langle (K - \langle K \rangle)^2 \rangle = \text{tr}(\sigma(K - \langle K \rangle)^2) \quad \text{with} \quad \langle K \rangle = \text{tr}(\sigma K). \quad (3.1)$$

But first we will compute its expectation value $\langle K \rangle$ and review the relationship between the boundary and bulk modular Hamiltonian.

3.1 The modular Hamiltonian: bulk versus boundary

We have defined the modular Hamiltonian K by the relation (1.8) in terms of the density matrix σ . This definition implies that the vacuum expectation $\langle K \rangle$ is non-zero. Indeed, one can easily show that $\langle K \rangle$ is equal to the entanglement entropy S ,

$$\langle K \rangle = \text{tr}(\sigma K) = -\text{tr}(\sigma \log \sigma) = S. \quad (3.2)$$

Together with the RT formula we may thus conclude that for a CFT with an AdS dual described by Einstein gravity that

$$\langle K \rangle = \frac{A(\Sigma)}{4G}. \quad (3.3)$$

As a next step we introduce a new operator ΔK that is defined as the deviation of the modular Hamiltonian K from its vacuum expectation value,

$$\Delta K \equiv K - \langle K \rangle \cdot \mathbf{1}. \quad (3.4)$$

This operator is well behaved in the sense that it has no divergences, and its vacuum expectation value is by definition equal to zero. With the above identifications, the definition of ΔK can be alternatively written as

$$K = \frac{A(\Sigma)}{4G} \cdot \mathbf{1} + \Delta K. \quad (3.5)$$

A noteworthy property of the operator ΔK is that its expectation value, when computed in a state ρ that differs from the vacuum state σ , is equal to the relative entropy. One has

$$\text{tr}(\rho \Delta K) = -\text{tr}(\rho \log \sigma) + \text{tr}(\sigma \log \sigma) \equiv S_{\text{rel}}(\rho|\sigma). \quad (3.6)$$

This result is familiar from ref. [13], and reflects the fact that the boundary relative entropy equals the bulk relative entropy. Furthermore, according to [13], the quantity ΔK represents the canonical energy of the bulk theory, which takes the form of the bulk modular Hamiltonian

$$\Delta K = \int_{\mathcal{C}} \xi_K^\mu T_{\mu\nu}^{\text{bulk}} d\mathcal{C}^\nu, \quad (3.7)$$

where \mathcal{C} is the volume denoted by the shaded area in figure 1. It is clear from this expression that ΔK sources the gravitational field in the bulk. In other words, it gravitates. Interestingly, it is also the same quantity that appears in the two-point function for the vacuum fluctuations of the modular Hamiltonian:

$$\langle \Delta K^2 \rangle = \langle (K - \langle K \rangle)^2 \rangle. \quad (3.8)$$

Our goal is to determine these fluctuations by applying the AdS/CFT dictionary. As as first step we note that, by inserting the definition eq. (1.8) of K , one can write the right-hand-side of eq. (3.8) explicitly in terms of the state σ as

$$\langle \Delta K^2 \rangle = \text{tr}(\sigma (\log \sigma)^2) - \left(\text{tr}(\sigma \log \sigma) \right)^2. \quad (3.9)$$

This equation manifestly shows that all the necessary information to compute $\langle \Delta K^2 \rangle$ is contained in the state σ itself. We will follow this path in the next subsection, where we apply the gravitational version of the replica method in which only the state σ is used as input.

3.2 The gravitational replica method

We will calculate the expectation value and the fluctuations in the modular Hamiltonian K by making use of a gravitational version of the replica method developed in [3–5]. Instead of introducing a finite inverse temperature β , we define free energies F_n ,

$$F_n = -\frac{1}{n} \log \left(\text{tr}(\sigma^n) \right), \quad (3.10)$$

where n is first chosen to take integer values.² One can think of the free energies F_n as being associated to thermal states ρ_n with inverse temperature equal to n times that of the vacuum state σ ,

$$\rho_n = \frac{\sigma^n}{\text{tr}(\sigma^n)} = \frac{e^{-nK}}{Z_n} \quad \text{with} \quad Z_n = \text{tr}(e^{-nK}) = e^{-nF_n}. \quad (3.11)$$

The partition function Z_n is related to the free energy F_n in the standard fashion.

The purpose of this section is to calculate the free energies F_n in AdS/CFT. Fortunately, this calculation was already performed in refs. [4, 5] by extending the method of ref. [3]. We will review these computations here: it will produce an expression for F_n for all integer values of n . Afterwards we will formally treat n as a continuum variable, and view F_n as an analytic function of n . This allows us to subsequently take its derivatives with respect to n . As we will explain below, the expectation value and fluctuations in K can be expressed in terms of these derivatives.

The gravitational replica method developed in refs. [3–5] amounts to considering the path integral over n -fold covers of an asymptotically AdS-Rindler spacetime. The gravitational path integral can be evaluated in the saddle point approximation, where the dominant saddle point represents a “cosmic brane” solution that is branched over the locus of the Rindler horizon [5]. In this way one finds that the free energy F_n can be expressed in terms of the classical gravitational action I_n for the cosmic brane,

$$nF_n = I_n - nI_1. \quad (3.12)$$

By requiring that the saddle point solution remains regular at the branch point, one learns that the underlying geometry of the base space, which is covered n -times by the full geometry, has an angle deficit equal to

$$\Delta\phi_n = 2\pi \left(1 - \frac{1}{n}\right). \quad (3.13)$$

The Lorentzian geometry of a cosmic brane with angle deficit (3.13) takes the form of the topological black hole metric eq. (2.10), where the blackening function is given by

$$f_n(r) = \frac{r^2}{L^2} - 1 - 2\Phi_n \frac{L^{d-2}}{r^{d-2}}. \quad (3.14)$$

As explained in the discussion around eq. (2.12), the quantity Φ_n is related to the tension (or energy density) of the cosmic brane. In this situation the required cosmic branes turn out to have negative tension.

The horizon of this geometry is located at $r = r_n$, where r_n is the solution to

$$f_n(r_n) = 0. \quad (3.15)$$

We will denote the co-dimension-two surface given by the bifurcate horizon of the cosmic brane by Σ_n . For $n = 1$ it coincides with the original Rindler-AdS horizon Σ . The inverse

²The free energies F_n are closely related to the Rényi entropies $S_n^{\text{Rényi}} = \frac{1}{1-n} \text{tr} \log \sigma^n$.

temperature T_n of the cosmic brane solution (or topological black hole) is given by

$$T_n = \frac{f'_n(r_n)}{4\pi}. \quad (3.16)$$

Since there is no cosmic brane for the case $n = 1$, we know that $\Phi_1 = 0$: in this case one finds $T_1 = 1/2\pi L$. By imposing that the angle deficit of the cosmic brane takes the value $\Delta\phi_n$ given in eq. (3.13), one deduces that the temperature T_n obeys

$$T_n = \frac{1}{2\pi L n}. \quad (3.17)$$

This equation, together with equation (3.15), forms a coupled set of equations from which both the radius r_n as well as the value of Φ_n can be determined. These equations are most conveniently written in terms of the dimensionless numbers $x_n = r_n/L$, where normalization is chosen so that $x_1 = 1$. By requiring that the inverse temperature T_n defined in eq. (3.16) gives the result eq. (3.17), one learns that the variables x_n are given by the positive solutions to the quadratic equation,

$$d x_n^2 - \frac{2}{n} x_n - (d-2) = 0 \quad \text{with} \quad x_n = r_n/L. \quad (3.18)$$

It is a simple exercise to obtain the solution to this equation and write an explicit expression for x_n in terms of n and d . Since the result will not be particularly useful for the following discussion, we leave this exercise to the reader. Instead we will use x_n itself to express the various other physical quantities. For instance, one can compute the value of Φ_n by imposing the condition that $f(r_n) = 0$. This leads to the result

$$\Phi_n = \frac{1}{2} (x_n^2 - 1) x_n^{d-2}. \quad (3.19)$$

This relation can be used to subsequently eliminate Φ_n from the remaining equations, so that all variables are expressed in terms of x_n .

It should be noted that all positive solutions to eq. (3.18) are less than one and only equal to one when $n = 1$. Hence, $x_n < 1$ for $n > 1$. This implies that Φ_n is negative, which also means that the “cosmic branes” described by the topological black hole geometries eqs. (2.10) and (3.14) have a negative tension and hence negative mass M .

Our next goal is to calculate the free energies F_n . In principle, this can be done by evaluating the classical action of the cosmic brane solutions. It appears, however, that there is a more direct and simple approach. Instead of directly calculating the Euclidean action I_n of the cosmic brane solutions, we follow the method of ref. [4]. By applying the familiar thermodynamic relation for the free energy $F(T) = E - TS$ together with the first law $dE = TdS$ one can derive the following integral expression for the free energy³

$$nF_n = \frac{1}{T(x_n)} \left[-T(x)S(x) \Big|_1^{x_n} + \int_1^{x_n} T(x)dS(x) \right], \quad (3.20)$$

³The dimensionless free energies F_n are related to the conventional free energy $F(T)$ via $nF_n = F(T_n)/T_n$.

where the functions

$$S(x) = \frac{A(\Sigma)}{4G} x^{d-1} \quad \text{and} \quad T(x) = \frac{1}{4\pi L} \left(dx - \frac{d-2}{x} \right) \quad (3.21)$$

represent the entropy and temperature associated with the horizon at $r/L = x$. The integral in this equation is easily evaluated, and together with the first term gives the following result for the free energy:

$$F_n = \frac{A(\Sigma)}{4G} \left(1 - \frac{1}{2} x_n^d - \frac{1}{2} x_n^{d-2} \right). \quad (3.22)$$

With this result it now becomes straightforward to compute the fluctuations in the modular Hamiltonian K .

3.3 Computation of the fluctuations in the modular Hamiltonian

The vacuum fluctuations of the modular Hamiltonian K as well as its vacuum expectation value can both be expressed as derivatives of the free energy F_n . First of all, the vacuum expectation value equals

$$\langle K \rangle = \frac{d}{dn} (n F_n) \Big|_{n=1} \quad (3.23)$$

One easily verifies this relation with the help of eq. (1.8). In a similar way one obtains the fluctuations in K by differentiating twice with respect to n . One has

$$\langle \Delta K^2 \rangle = - \frac{d^2}{dn^2} (n F_n) \Big|_{n=1} \quad (3.24)$$

To check this last relation one inserts eq. (1.8) into the left-hand-side, or uses the definition eq. (3.10) for F_n on the right-hand-side.

To evaluate these derivatives of the free energy F_n with respect to n , we use its expression eq. (3.22) in terms of the variables x_n . First let us determine the expectation value of the modular energy K in the infinite set of generalized thermal states ρ_n defined in eq. (3.11). These expectation values are given in terms of the free energy F_n by

$$\langle K \rangle_n \equiv \text{tr}(K \rho_n) = \frac{\text{tr}(K \sigma^n)}{\text{tr}(\sigma^n)} = \frac{d}{dn} (n F_n). \quad (3.25)$$

Here we keep n free. To compute $\langle K \rangle_n$ for general values of n we need to know the derivative of x_n with respect to n . By multiplying the quadratic relation eq. (3.18) by $n/2x_n$ and differentiating with respect to n one finds:

$$n \left(dx_n + \frac{d-2}{x_n} \right) \frac{dx_n}{dn} = -dx_n + \frac{d-2}{x_n}. \quad (3.26)$$

With the help of this relation one can now easily compute the vacuum expectation value of K and its vacuum fluctuations. First of all, for the expectation value $\langle K \rangle_n$ we find

$$\langle K \rangle_n = \frac{A(\Sigma)}{4G} \left(1 + \frac{d-1}{2} (x_n^2 - 1) x_n^{d-2} \right). \quad (3.27)$$

By taking $n = 1$ one verifies that the vacuum expectation value of K is, as expected, equal to the entanglement entropy

$$\langle K \rangle = \frac{A(\Sigma)}{4G}. \quad (3.28)$$

The next step is to evaluate the fluctuations. Using the representation (3.22) it is straightforward to show that

$$\langle \Delta K^2 \rangle = -\frac{d^2}{dn^2} \left(n F_n \right) \Big|_{n=1} = -\frac{d}{dn} \langle K \rangle_n \Big|_{n=1} = -\frac{1}{n} \frac{d}{dn} S(x_n) \Big|_{n=1}, \quad (3.29)$$

where in the last step we made use of the first law of thermodynamics. This last expression can also be directly derived by differentiating the integral expression (3.20) twice with respect to n and make use of the fact that $1/T(x_n) = 2\pi L n$. The right hand side can be evaluated in a straightforward way and gives

$$\frac{d}{dn} S(x_n) \Big|_{n=1} = (d-1) \frac{A(\Sigma)}{4G} x_n^{d-2} \frac{dx_n}{dn} \Big|_{n=1} = -\frac{A(\Sigma)}{4G}, \quad (3.30)$$

where the derivative of x_n is evaluated via the relation (3.26): $dx_n/dn|_{n=1} = -1/(d-1)$. In this way we arrive at the final result for the fluctuations in the modular Hamiltonian⁴

$$\langle \Delta K^2 \rangle = \frac{A(\Sigma)}{4G}. \quad (3.31)$$

This is indeed the answer that we anticipated in [1], and can be argued for on general grounds, since it expresses the fact that the size of the fluctuations $\sqrt{\langle \Delta K^2 \rangle}$ obeys the familiar “square root” law for a gaussian stochastic process.

4 Backreaction due to the fluctuating modular energy

After having established that the modular Hamiltonian fluctuates in the vacuum, we now come to the question of how these fluctuations influence the geometry. As we made clear, the vacuum expectation value of the modular Hamiltonian does not gravitate, since this would imply that the vacuum expectation value of the geometry would not be equal to that of the vacuum. The difference ΔK between the value of the modular Hamiltonian K and its vacuum expectation value does gravitate. In particular, as we will show, a state for which the expectation value of K differs from that of the vacuum, will correspond to a geometry different from empty AdS.

Specifically we will show that a non-zero expectation value of ΔK results in a non-zero (expectation) value of the parameter Φ that we identified with the Newtonian potential. To show this fact, we will first relate ΔK and the mass M that appears in eq. (2.12) for Φ . Subsequently we will argue that the same relation between ΔK and Φ also holds at the level of the fluctuations. This implies in particular that Φ fluctuates and that as a result the location of the horizon exhibits vacuum fluctuations. Our reasoning in this section is inspired by [14] and will closely follow the discussion in [1].

⁴After completing our work we learned of related papers in a similar result was obtained, see *Note added*.

4.1 The relation between the mass M and the modular hamiltonian K

The aim of this subsection is to show that the mass M , as defined by eq. (2.12), can, as a quantum operator, be identified with

$$M = \frac{1}{2\pi L} (K - \langle K \rangle). \quad (4.1)$$

Here the pre-factor $1/2\pi L$ represents the Hawking temperature at the Rindler horizon. In other words, the mass operator M is, up to an overall factor given by the temperature, equal to difference of the modular Hamiltonian K and its vacuum expectation value $\langle K \rangle$. A crucial ingredient in the derivation of this relation is the fact that the mass M satisfies the first law of black hole thermodynamics

$$dM = TdS. \quad (4.2)$$

Our proof of eq. (4.1) amounts to showing that both sides of this equation obey the same 1st law of black hole thermodynamics.

First, let us derive that the mass M , as defined by eq. (2.11) and eq. (2.12), indeed obeys the first law given in eq. (4.2). For this we use the fact that the location of the horizon is determined by the equation $f(r_h, M) = 0$ for all values of M . Hence, the variation dr_h of the horizon location and the mass variation dM are related by

$$df(r_h, M) = f'(r_h, M)dr_h + \frac{\partial f(r_h, M)}{\partial M}dM = 0. \quad (4.3)$$

From the expressions in eq. (2.11), (2.12) we learn that

$$\frac{\partial f(r_h, M)}{\partial M} = -\frac{16\pi G}{(d-1)V_{d-1}r_h^{d-2}}. \quad (4.4)$$

It is now a simple exercise to show, with the help of eq. (3.16) and by varying the area-entropy relation in eq. (2.7), that the variation dM indeed satisfies the first law of black hole thermodynamics given in eq. (4.2).

Our next goal is to show that the right-hand side of eq. (4.1) obeys the same first law of black hole entropy, when K is defined by eq. (1.8). We will do this at the level of expectation values in the infinite set of states ρ_n defined in eq. (3.11). For this purpose it will be useful to rewrite eq. (4.2) in terms of the free-energy $F(T) = M - TS$ in the familiar form

$$dF = -SdT \quad (4.5)$$

To show that this relation also holds for the right-hand side of eq. (4.1) we consider the expectation value in the infinite set of states ρ_n defined in eq. (3.11). First of all, one easily verifies that the dimensionless free energies F_n defined in eq. (3.10) obey

$$F_n \equiv -\frac{1}{n} \log \text{tr}(\sigma^n) = \langle K \rangle_n - \frac{1}{n} S_n, \quad (4.6)$$

where $S_n = -\text{tr}(\rho_n \log \rho_n)$ is the von Neumann entropy⁵ associated with the state ρ_n . By taking the differential of the first expression one derives that F_n satisfies

$$dF_n = S_n \frac{dn}{n^2} \quad (4.7)$$

which is the direct analogue of eq. (4.5). In these steps $1/n$ plays the role of a dimensionless temperature, and is considered to be a continuous variable. This can be made even more explicit by rewriting this relation, using the second expression for F_n in eq. (3.22), in the form

$$d\langle K \rangle_n = \frac{1}{n} dS_n. \quad (4.8)$$

This is the direct analogue of the first law in eq. (4.2). Now note that the physical temperature T_n defined in (3.17) differs from $1/n$ by a factor $2\pi L$. The same factor appears therefore in the relation between the mass M and the modular Hamiltonian K . Finally, by requiring that the mass M , defined as an operator, has a zero vacuum expectation value, one uniquely fixes the relation between the mass and the modular Hamiltonian to take the form eq. (4.1).

As a final check on this identification, we note that inside the causal diamond on the boundary the modular operator K acts as the generator of a conformal boost in the CFT, while in the bulk it represents a Lorentz boost that leaves the Rindler horizon invariant. The corresponding (conformal) Killing vector ξ_K is related to the generator $\partial/\partial t$ of translation in AdS-Rindler time by a factor equal to the temperature associated with the Rindler horizon. Hence, we have

$$\xi_K^\mu \partial_\mu = 2\pi L \frac{\partial}{\partial t}. \quad (4.9)$$

This equation again implies that the factor $2\pi L$ must also appear in the relation between the mass M and the quantity $K - \langle K \rangle$.

4.2 Fluctuations in the Newtonian potential on the Rindler horizon

In this subsection we take the next step to show that the fluctuations in the modular Hamiltonian associated with the causal region also leads to metric fluctuations, specifically in the Newtonian potential Φ . By inserting this relation into the expression eq. (2.12) we arrive at the expression for Φ ,

$$\Phi = \frac{\Delta K}{(d-1)} \frac{4G}{A(\Sigma)} \quad \text{with} \quad \Delta K = K - \langle K \rangle. \quad (4.10)$$

As announced, this relation shows that ΔK gravitates. Since ΔK has a zero vacuum expectation value, it induces no change in the expectation value of the metric, but the fact that ΔK exhibits vacuum fluctuations means that it can lead to small quantum fluctuations in the value of the Newton potential Φ on the AdS-Rindler horizon.

Just as in our discussion for the mass M we will again read this equation as an operator identity relating Φ to ΔK . In making this identification, we will regard the area

⁵Note that the entropies S_n differ from the Rényi entropies $S_n^{\text{Rényi}}$ defined in the footnote below eq. (3.10).

of the Rindler horizon $A(\Sigma)$ as a scalar quantity that is fixed: it represents the size of the Hilbert space and the entanglement entropy of the vacuum state. As such it is not a quantum operator.

To further justify the relation eq. (4.10) we will first verify it again at the level of expectation values in the state ρ_n . We want to verify that the expectation value $\langle \Phi \rangle_n$ of Φ in the state ρ_n is given by the classical result that we determined in eq. (3.19). In other words, we want to show that we can take $\langle \Phi \rangle_n \equiv \Phi_n$. To verify this relation we compute the expectation value of the right-hand side of eq. (4.10). This gives

$$\langle \Phi \rangle_n = \frac{\langle \Delta K \rangle_n}{(d-1)} \frac{4G}{A(\Sigma)}. \quad (4.11)$$

By comparing the expressions (3.19) and (3.27) for Φ_n and $\langle K \rangle_n$, and after subtracting the vacuum value eq. (3.28) for $\langle K \rangle$, one easily checks that the right-hand side is equal to Φ_n , as required. This therefore suggests that eq. (4.10) also holds at the level of operators.

If we assume that this operator identification indeed holds, we can now use it to compute the fluctuations in the Newton potential Φ in a simple and straightforward way. We find that the value of Φ on the horizon has a non-zero two-point function in the vacuum state σ given by

$$\langle \Phi^2 \rangle = \frac{\langle \Delta K^2 \rangle}{(d-1)^2} \left(\frac{4G}{A(\Sigma)} \right)^2 = \frac{1}{(d-1)^2} \frac{4G}{A(\Sigma)}, \quad (4.12)$$

where in the last step we used our result eq. (3.31) for the fluctuations in ΔK .

After having argued that the fluctuations in the modular energy ΔK also source fluctuations in the geometry, in particular in the Newtonian potential Φ on the horizon, we will now investigate the consequences of such fluctuations.

4.3 Horizon location and photon travel time

Let us return to the AdS spacetime in Poincaré coordinates

$$ds^2 = \frac{L^2}{z^2} (dz^2 + dx_i^2 - dx_0^2), \quad (4.13)$$

and again consider the causal domain bounded by the AdS-Rindler horizon. The horizon can be thought of as being made up of many photon trajectories that are simultaneously emitted from the boundary, and subsequently reflect off an (imaginary) mirror located on the bifurcate horizon. Our aim in this subsection is to use this observation to show that the travel time of a real photon, emitted from the boundary and reflected off a real mirror, is directly correlated with the location of the associated Rindler horizon.

The coordinate transformation from the Poincaré coordinates to the Rindler coordinates given in eq. (2.5) implies the following relation

$$\left(\frac{(R - x_0)^2 - z^2 - x_i^2}{2Rz} \right) \left(\frac{(R + x_0)^2 - z^2 - x_i^2}{2Rz} \right) = \frac{r^2}{L^2} - 1. \quad (4.14)$$

This equation expresses when we take $r = L$ that either the first or the second factor on the left-hand-side vanishes. The resulting equations describe the location of the lower

and upper parts of the causal diamond in the bulk. The intersection of the two equations corresponds to the bifurcate horizon, which is hence located at $x_0 = 0$ and $z^2 + x_i^2 = R^2$.

We now consider a photon trajectory which travels along $x_i = 0$ and reflects off a mirror located on the bifurcate horizon at $z = R$. We can describe its path by disregarding the x_i coordinates in eq. (4.14). In this way one finds that the photon is emitted and received at the boundary at $x_0 = -R$ and $x_0 = R$. These equations hold when the boundary is chosen to be located at $z = 0$. However, to regulate the horizon area and to obtain a finite induced metric on the boundary, we will now introduce a small cut-off z_c by restricting the range of the z -coordinate to

$$z \geq z_c, \quad \text{with} \quad z_c \ll R. \quad (4.15)$$

The AdS-boundary is thus now located at $z = z_c$. The emission, reflection and reception times of a photon, that is emitted from $z = z_c$ and again reflected from a mirror in the bulk at $z = R$, are now given by

$$x_0^{\text{emission}} = -R + z_c, \quad x_0^{\text{reflection}} = 0, \quad \text{and} \quad x_0^{\text{reception}} = R - z_c. \quad (4.16)$$

Hence, after introducing the cut-off, the round trip time of the photon equals $2(R - z_c)$: this is equal to twice the difference in the z -coordinate of the mirror and the boundary.

The *physical time* that is measured by a clock on the boundary at $z = z_c$ is, however, not equal to x_0 . Namely, due to the pre-factor $(L/z_c)^2$ in front of the metric, the actual time measured on a physical clock is given by Lx_0/z_c . The physical round trip time $T_{\text{r.t.}}$ of the photon is therefore,

$$T_{\text{r.t.}} = \frac{L}{z_c} \cdot 2(R - z_c). \quad (4.17)$$

Note that this round trip time would diverge if we send the cut-off $z_c \rightarrow 0$.

A second motivation for introducing the cut-off z_c is that the area $A(\Sigma)$ of the Rindler horizon remains finite. Previously we have shown that this area is given by eqs. (2.8) and (2.9) in terms of another cut-off χ_c . To determine the relationship between the two cut-offs z_c and χ_c we observe that the horizon is located at

$$z^2 + x_i^2 = R^2 \quad \text{with} \quad |x_i| = R \tanh \chi. \quad (4.18)$$

Imposing that $\chi \leq \chi_c$ leads to the inequalities

$$x_i^2 \leq R^2 \tanh^2 \chi_c \quad \text{hence} \quad z^2 \geq R^2 - R^2 \tanh^2 \chi_c. \quad (4.19)$$

From these equations we learn that the relationship between z_c and χ_c is given by

$$z_c = R \sqrt{1 - \tanh^2 \chi_c} = \frac{R}{\cosh \chi_c}. \quad (4.20)$$

In the following we will be interested in comparing the value of the photon round trip time with the horizon area $A(\Sigma)$. For this purpose we express both in terms of the same cut-off parameter χ_c . After inserting eq. (4.20) into eq. (4.17) we find for the round trip time

$$T_{\text{r.t.}} = 2L(\cosh \chi_c - 1). \quad (4.21)$$

For later convenience we also repeat here the full expression for the horizon area

$$A(\Sigma) = L^{d-1} \Omega_{d-2} \int_0^{\chi_c} d\chi \sinh^{d-2} \chi. \quad (4.22)$$

Both these equations hold in arbitrary dimensions. The discussion in this subsection has been completely classical, and has so far ignored any fluctuations in the geometry. The aim of the following subsection is to determine what happens to the location of the horizon and the photon travel times due to the fluctuations in the Newtonian potential Φ .

4.4 Fluctuations in horizon location and photon travel time

By introducing the cut-off χ_c we have regularized the horizon area $A(\Sigma)$, and turned it into a finite quantity. This implies that the fluctuations in the Newton potential Φ given in eq. (4.12) are non-vanishing. The purpose of this subsection is to discuss the physical implications of the fluctuations in Φ on the horizon location and the photon travel time.

Let us assume that quantum fluctuations cause Φ , at some moment in time, to acquire a small but non-zero value of the order of

$$\Phi \sim \sqrt{\langle \Phi^2 \rangle} = \frac{1}{d-1} \sqrt{\frac{4G}{A(\Sigma)}}. \quad (4.23)$$

In other words, we regard Φ as a stochastic variable, whose variance is given by $\langle \Phi^2 \rangle$. First let us determine the perturbation in the location of the Rindler horizon due to Φ . In terms of the Rindler-AdS coordinates, it is located at the solution of $f(r_h) = 0$. Assuming that $\Phi \ll 1$, we find that the perturbed horizon is now located at

$$\frac{r_h^2}{L^2} - 1 = \Phi. \quad (4.24)$$

The fluctuations in Φ thus induce a quantum uncertainty in the location of the horizon. This result is in accordance with the arguments of ref. [14]. It will be useful to rewrite the perturbed location of the horizon given in eq. (4.24) in terms of Poincaré coordinates. Inserting eq. (4.24) into eq. (4.14) gives

$$\left(\frac{(R - x_0)^2 - z^2 - x_i^2}{2Rz} \right) \left(\frac{(R + x_0)^2 - z^2 - x_i^2}{2Rz} \right) = 2\Phi. \quad (4.25)$$

Our next goal is to compute the effect on the travel time of the photon. For this purpose we will now make the physical assumption that the uncertainty in the location of the reflection point of the photon is equal to the uncertainty in the location of the bifurcate horizon. We will again choose the reflection moment with the mirror to be at $x_0 = 0$ and to be located at $x_i = 0$. Hence, we are only considering fluctuations in the z -coordinate. One easily works out that the solution of eq. (4.25) for the coordinate z , when we take $x_0 = x_i = 0$, is given by

$$\frac{R^2 - z^2}{2Rz} = \pm \sqrt{2\Phi} \quad \implies \quad z^{\text{reflection}} = R \left(\sqrt{1 + 2\Phi} \pm \sqrt{2\Phi} \right). \quad (4.26)$$

Hence the quantum uncertainty in location of the reflection point is $\Delta z^{\text{reflection}} = R\sqrt{2\Phi}$, where we ignored terms of order Φ . Note that we could have also phrased this calculation as the reflection point in the bulk remaining fixed, but the size of the causal diamond on the boundary shifting from $R \rightarrow \tilde{R} = R(\sqrt{1+2\Phi} \pm \sqrt{2\Phi})$. Either calculation gives the same time delay. This is because the physical result can be found equally as a (passive or active) coordinate transformation on the boundary or in the bulk.

As we explained, the time it takes in terms of x_0 for a photon to traverse between the mirror and the boundary is equal to the difference in the z -coordinates of the mirror and the boundary. This implies that the quantum uncertainty in the z -coordinate directly translates into a quantum uncertainty in the emission and reception times. We may thus conclude that

$$\left. \frac{\Delta z^2}{z^2} \right|_{\text{reflection}} = \left. \frac{\Delta x_0^2}{x_0^2} \right|_{\text{emission}} = \left. \frac{\Delta x_0^2}{x_0^2} \right|_{\text{reception}} = 2\Phi. \quad (4.27)$$

For the physical round trip time for the photon we reach a similar conclusion. In fact, the ratio of the fluctuations $\Delta T_{\text{r.t.}}$ to the travel time $T_{\text{r.t.}}$ is independent of the metric pre-factor and hence is given in terms of Φ by a similar expression as eq. (4.27).

The statements made thus far involve a single photon trajectory. Observables, however, involve interference of light following two different trajectories. Further, $\langle \Phi \rangle = 0$, such that one must consider the *variance* of eq. (4.27). Thus the first contribution to the observable enters as a four-point correlation function of the observable:

$$\left\langle \left(\frac{\Delta T_{\text{r.t.}}^2}{T_{\text{r.t.}}^2} \right)^2 \right\rangle = 4\langle \Phi^2 \rangle. \quad (4.28)$$

In an experiment measuring the phase of light, $\frac{\Delta T_{\text{r.t.}}^2}{T_{\text{r.t.}}^2}$ is correlated over two different paths. Making use of eq. (4.23), we thus have

$$\frac{\Delta T_{\text{r.t.}}^2}{T_{\text{r.t.}}^2} = \frac{2}{d-1} \sqrt{\frac{4G}{A(\Sigma)}}. \quad (4.29)$$

Thus we see, because the first non-zero contribution to fluctuations in time-of-arrival of a pulse of light enters at the four-point, and because the gravitational potential, $\langle \Phi^2 \rangle \sim \frac{G^2}{L^2} S$ is enhanced by a factor of the entropy (containing $S \sim 1/G$), the fluctuations in the observable scale with \sqrt{G} .

We now compare this surprising result to that obtained in our previous work ref. [1] for $4d$ Minkowski space. For this purpose we take the dimensions to be $d = 3$, so that we are dealing with a $4d$ AdS spacetime. The expression in eq. (4.22) for the area $A(\Sigma)$ is in this case easily evaluated,

$$A(\Sigma) \stackrel{d=3}{=} 2\pi L^2 (\cosh \chi_c - 1). \quad (4.30)$$

By inserting this into our general result, eq. (4.29), we find for the fluctuations in the photon travel time

$$\frac{\Delta T_{\text{r.t.}}^2}{T_{\text{r.t.}}^2} \stackrel{d=3}{=} \frac{l_p}{2\pi L} \cdot \frac{1}{\sqrt{\cosh \chi_c - 1}}, \quad (4.31)$$

where we identified the $4d$ Planck scale with $l_p^2 = 8\pi G$. We thus confirm the result of ref. [1], that the fluctuations in the photon travel times scale like $\sqrt{l_p}$, with a denominator that is determined by the light crossing time. The difference in the denominator of this expression, relative to our previous result in flat space, is due the negative curvature of AdS space. Also note that in ref. [1], the fluctuations in eq. (4.28) were correlated over two different paths, with a correlation strength that (in flat four-dimensional space) is given by spherical harmonics.

Lastly, we note that the flat space results in ref. [1] can be reconstructed in a Randall-Sundrum II scenario, where the boundary is pulled into the bulk, inducing gravity on the brane. When $\chi_c \gg 1$, we have

$$\frac{\Delta T_{\text{r.t.}}^2}{T_{\text{r.t.}}^2} = \frac{2^{d/2}}{d-1} \sqrt{\frac{4(d-2)G_{\text{bulk}}}{\Omega_{d-2}L^{d-1}e^{(d-2)\chi_c}}} = \frac{2^{d/2}}{d-1} \sqrt{\frac{4G_{\text{bdy}}}{\Omega_{d-2}L^{d-2}e^{(d-2)\chi_c}}} \stackrel{d=4}{\approx} \sqrt{\frac{G_{\text{bdy}}}{T_{\text{r.t.}}^2}}, \quad (4.32)$$

where we have used the fact that the bulk and boundary gravitational constants are related by $G_{\text{bdy}} = (d-2)G_{\text{bulk}}/L$. We intend to pursue this model further in future work.

5 Discussion

The main aim of this work was to study vacuum energy fluctuations in AdS space that occur in a finite spacetime region, and to study their implications for the spacetime geometry. In particular we analyzed their effect on the travel times of photons that reflect off a mirror in the bulk. An important step in our argumentation was that, for the problem of interest, one may focus on the causal diamond associated with a single round trip of a photon. We showed that, while the expectation value in the energy remains zero, there exist local energy fluctuations of the size given by eq. (3.31) in any finite region of the AdS-spacetime. It is important to point out, however, that the relative size of these local energy fluctuations, and their effect on the geometry, goes to zero when we take the size of the region to infinity. We have shown that for a finite region, the energy fluctuations source metric fluctuations in the bulk, given by eq. (4.10). We subsequently showed that a light beam reflecting off a mirror in the bulk experiences fluctuations in the time it leaves and arrives from the boundary, summarized in eq. (4.29). We again emphasize that these effects only occur because one is measuring a finite part of the spacetime.

While our results in AdS/CFT are in qualitative agreement with those obtained in Minkowski space in ref. [1], in the Minkowski case one has less control over the underlying microscopics. In ref. [1], we only presented a heuristic argument for the size of the fluctuations given by eq. (3.31), where in the $4d$ flat space situation the surface Σ has an area $A = 4\pi R^2$ and divides the inside and outside of the causal diamond with finite radius R . In this paper we verified these arguments in the context of AdS/CFT, and provided further evidence for the assumptions that were made for the flat space case. In particular, one would like to understand to what extent flat space holographic assumptions, like the ones presented in this paper, are justified.

There are many interesting directions to pursue to clarify this question. First, moving the boundary CFT into the bulk, as in a braneworld scenario, one could study energy

and metric fluctuations on the brane with Minkowski metric. Second, $\langle \Delta K^2 \rangle$ could be computed directly with the quantized Einstein-Hilbert metric. We expect that such a calculation would yield energy fluctuations proportional to the surface area (dividing the inside and out of the causal region) but quadratically UV divergent, similar to the classic calculation of the entanglement entropy [15, 16].

Ultimately we are seeking to show that, at least where certain soft properties of the quantized Einstein-Hilbert action are concerned, the holographic principle applies equally well in Minkowski space as in the context of AdS/CFT. There has already been progress in these directions in the context of soft theorems, see for example [17–20]. The finite causal diamond and its realization in the experimental set-up of an interferometer provides a novel context in which to pursue these questions further.

Acknowledgments

We thank Nima Arkani-Hamed, Tom Banks, Dan Carney, Cliff Cheung, David Gross, Cynthia Keeler, Takemichi Okui and Raman Sundrum for discussions, and Eric Perlmutter, Jan de Boer and for bringing ref. [8], [9], and [10] to our attention.

Open Access. This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY 4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] E.P. Verlinde and K.M. Zurek, *Observational Signatures of Quantum Gravity in Interferometers*, [arXiv:1902.08207](https://arxiv.org/abs/1902.08207) [[INSPIRE](#)].
- [2] H. Casini, M. Huerta and R.C. Myers, *Towards a derivation of holographic entanglement entropy*, *JHEP* **05** (2011) 036 [[arXiv:1102.0440](https://arxiv.org/abs/1102.0440)] [[INSPIRE](#)].
- [3] A. Lewkowycz and J. Maldacena, *Generalized gravitational entropy*, *JHEP* **08** (2013) 090 [[arXiv:1304.4926](https://arxiv.org/abs/1304.4926)] [[INSPIRE](#)].
- [4] L.-Y. Hung, R.C. Myers, M. Smolkin and A. Yale, *Holographic Calculations of Renyi Entropy*, *JHEP* **12** (2011) 047 [[arXiv:1110.1084](https://arxiv.org/abs/1110.1084)] [[INSPIRE](#)].
- [5] X. Dong, *The Gravity Dual of Renyi Entropy*, *Nature Commun.* **7** (2016) 12472 [[arXiv:1601.06788](https://arxiv.org/abs/1601.06788)] [[INSPIRE](#)].
- [6] S. Ryu and T. Takayanagi, *Holographic derivation of entanglement entropy from AdS/CFT*, *Phys. Rev. Lett.* **96** (2006) 181602 [[hep-th/0603001](https://arxiv.org/abs/hep-th/0603001)] [[INSPIRE](#)].
- [7] V.E. Hubeny, M. Rangamani and T. Takayanagi, *A covariant holographic entanglement entropy proposal*, *JHEP* **07** (2007) 062 [[arXiv:0705.0016](https://arxiv.org/abs/0705.0016)] [[INSPIRE](#)].
- [8] E. Perlmutter, *A universal feature of CFT Rényi entropy*, *JHEP* **03** (2014) 117 [[arXiv:1308.1083](https://arxiv.org/abs/1308.1083)] [[INSPIRE](#)].
- [9] Y. Nakaguchi and T. Nishioka, *A holographic proof of Rényi entropic inequalities*, *JHEP* **12** (2016) 129 [[arXiv:1606.08443](https://arxiv.org/abs/1606.08443)] [[INSPIRE](#)].

- [10] J. De Boer, J. Järvelä and E. Keski-Vakkuri, *Aspects of capacity of entanglement*, *Phys. Rev. D* **99** (2019) 066012 [[arXiv:1807.07357](#)] [[INSPIRE](#)].
- [11] T. Faulkner, M. Guica, T. Hartman, R.C. Myers and M. Van Raamsdonk, *Gravitation from Entanglement in Holographic CFTs*, *JHEP* **03** (2014) 051 [[arXiv:1312.7856](#)] [[INSPIRE](#)].
- [12] T. Jacobson, *Entanglement Equilibrium and the Einstein Equation*, *Phys. Rev. Lett.* **116** (2016) 201101 [[arXiv:1505.04753](#)] [[INSPIRE](#)].
- [13] D.L. Jafferis, A. Lewkowycz, J. Maldacena and S.J. Suh, *Relative entropy equals bulk relative entropy*, *JHEP* **06** (2016) 004 [[arXiv:1512.06431](#)] [[INSPIRE](#)].
- [14] D. Marolf, *On the quantum width of a black hole horizon*, *Springer Proc. Phys.* **98** (2005) 99 [[hep-th/0312059](#)] [[INSPIRE](#)].
- [15] M. Srednicki, *Entropy and area*, *Phys. Rev. Lett.* **71** (1993) 666 [[hep-th/9303048](#)] [[INSPIRE](#)].
- [16] S.N. Solodukhin, *Entanglement entropy of black holes*, *Living Rev. Rel.* **14** (2011) 8 [[arXiv:1104.3712](#)] [[INSPIRE](#)].
- [17] D. Kapec, P. Mitra, A.-M. Raclariu and A. Strominger, *2D Stress Tensor for 4D Gravity*, *Phys. Rev. Lett.* **119** (2017) 121601 [[arXiv:1609.00282](#)] [[INSPIRE](#)].
- [18] D. Kapec, A.-M. Raclariu and A. Strominger, *Area, Entanglement Entropy and Supertranslations at Null Infinity*, *Class. Quant. Grav.* **34** (2017) 165007 [[arXiv:1603.07706](#)] [[INSPIRE](#)].
- [19] C. Cheung, A. de la Fuente and R. Sundrum, *4D scattering amplitudes and asymptotic symmetries from 2D CFT*, *JHEP* **01** (2017) 112 [[arXiv:1609.00732](#)] [[INSPIRE](#)].
- [20] A. Ball, E. Himwich, S.A. Narayanan, S. Pasterski and A. Strominger, *Uplifting AdS_3/CFT_2 to flat space holography*, *JHEP* **08** (2019) 168 [[arXiv:1905.09809](#)] [[INSPIRE](#)].